

Mild whipworm infection diagnosed by doing colonoscopy in our study

Govindarajalu Ganesan

Associate professor, Dept. of General surgery, Indira Gandhi Medical College and Research Institute, Vazhudavoor Rd, Kathirkamam, Puducherry, India

Abstract

Objective: To find the grade of whipworm infection diagnosed by doing colonoscopy in our study.

Methods: A study of 72 patients who had undergone colonoscopy for a period of 5 years from November 2009 to October 2014 was carried out. In patients found to have parasitic worms during colonoscopy, number of parasitic worms in the colon were noted and investigations were done to know about the presence or absence of anaemia to know about the severity and grade of parasitic infection.

Results: Out of these 72 patients, parasitic worm was found in the colon in only one patient. The parasitic worm found in this patient was identified as whipworm or trichuris trichiura by its characteristic whip like shape. It has a short posterior thick part resembling the short handle of the whip and a long, thin anterior part resembling the distal long, thin part of the whip. Our patient was found to have only single whipworm in the colon while doing colonoscopy. Our patient did not have anaemia (haemoglobin 14.4 g%) and did not have any serious complications like Trichuris dysentery syndrome causing anaemia or colonic obstruction and perforation.

Conclusion: Hence our patient was found to have only mild whipworm infection with only a single whipworm in the colon while doing colonoscopy without anaemia and without serious complications like Trichuris dysentery syndrome or colonic obstruction and perforation.

Keywords: adult whipworm, trichuris trichiura, mild whipworm infection, colonoscopy

Introduction

Whipworms are the most common nematodes or roundworms found in the large intestine of human beings while doing colonoscopy. There has been also reports of finding whipworm in the large intestine of human beings while doing colonoscopy in many parts of the world [1-14]. In almost all the studies, whipworm or trichuris trichiura was almost the only intestinal helminth or roundworm found in the large intestine of human beings while doing colonoscopy [2]. Our patient was also found to have whipworm in the colon while doing colonoscopy.

Materials and Methods

This study was conducted in the department of general surgery, Aarupadai Veedu Medical College and Hospital, Puducherry. A study of 72 patients who had undergone colonoscopy for a period of 5 years from November 2009 to October 2014 was carried out. In patients found to have parasitic worms during colonoscopy, number of parasitic worms in the colon were noted and investigations were done to know about the presence or absence of anaemia to know about the severity and grade of parasitic infection. Anaemia is defined as haemoglobin < 12g/dl or 12g% in women and haemoglobin < 13g/dl or 13g% in men.

Results

Out of these 72 patients, parasitic worm was found in only one patient. The parasitic worm found in this patient was identified as whipworm or trichuris trichiura by its characteristic whip like shape. This patient was an eighty year old male patient and presented with history of lower

abdominal pain for 1 week, constipation for 2 days and vomiting for 1 day. On abdominal examination, his abdomen was soft, mildly distended and mild tenderness was present in the hypogastric region. His haemoglobin was 14.4g%. When he was subjected to diagnostic colonoscopy, one adult whipworm was found in the sigmoid colon while doing colonoscopy. The patient was treated with a single dose of 400mg of albendazole and started showing clinical improvement.

Absence of anaemia in our patient

In our patient with whipworm found in colonoscopy, anaemia was not present [(haemoglobin 14.4 g%)]. In another study also [5], all the three patients with whipworm or trichuris trichiura found in colonoscopy did not have anaemia (haemoglobin 13.7 g %, 14.1 g %, 13.9 g %).

Absence of anaemia in another study

In the study conducted by Kyung-Sun Ok *et al* [5] in Korea, colonoscopy revealed a small, white, worm-like object attached to the cecum (trichuris trichiura) in a 56 year old male patient without anaemia (haemoglobin 13.7 g %). In the same study [5], colonoscopy revealed a small, white, worm on the proximal ascending colon (trichuris trichiura) in a 46 year old male patient without anaemia (haemoglobin 14.1 g %). In the same study [5], colonoscopy revealed a small, white, worm within the ileocecal valve (trichuris trichiura) without anaemia (haemoglobin 13.9 g %) in a 55 year old male patient.

Mild whipworm infection in our patient

Our patient was found to have only mild whipworm infection

without anaemia (haemoglobin 14.4 g%) and without serious complications like Trichuris dysentery syndrome causing anaemia requiring intensive medical treatment and without colonic obstruction and perforation requiring surgical resection of the colon.

Mild whipworm infection in three patients in another study

In the study conducted by Kyung-Sun Ok *et al* [5] in Korea, all the three patients with whipworm or trichuris trichiura found in colonoscopy were also found to have only mild whipworm infection without anaemia (haemoglobin 13.7 g %, 14.1 g%, 13.9 g%) and without serious complications like Trichuris dysentery syndrome causing anaemia requiring intensive medical treatment and without colonic obstruction and perforation requiring surgical resection of the colon. All the three patients were treated with albendazole and were symptom-free after treatment with albendazole.

Discussion

i) Blood loss and anaemia in whipworm infection

The male whipworm is 3 to 4.5 cm and the female whipworm is 3.5 to 5 cm in length [11]. Unlike hookworm which sucks blood from the small intestinal mucosa, whipworm [(trichuris trichiura) does not suck blood and feeds only on the tissue secretions of the large intestinal wall. Also most people in endemic areas of Trichuris trichiura infections are colonized

only by a small number of worms (usually less than 15) unlike large number of hookworms present commonly in hookworm infection. Hence anaemia is not common in whipworm infection, but is common in hookworm infection. Our patient with whipworm infection also did not have any anaemia (haemoglobin 14.4 g%).

ii) Whitish colour of whipworm

Whipworms do not feed on blood and feeds only on the tissue secretions of the large intestinal wall. Whipworm is always whitish in colour [5] as it does not feed on blood (fig 1, 2).

iii) Only a very small portion of the long anterior part of whipworm seen during colonoscopy

We can see only the short posterior thick part of whipworm entirely in the lumen of the large intestine [13, 14] but only a very small portion of the long, thin anterior part while doing colonoscopy since most of the anterior part penetrates into the large intestinal wall in order to feed on the tissue secretions of the large intestinal wall [13, 14]. Hence in fig 1, we can see only the short posterior thick part of the whitish coloured whipworm entirely in the lumen of the sigmoid colon but only a very small portion of the long, thin anterior part since most of the anterior part penetrates into the large intestinal wall for feeding purpose. But in the highly magnified view in fig 2, we can see clearly both the short posterior thick part and also the anterior thin part clearly due to the higher magnification.




Fig 1: Showing clearly and entirely only the short, posterior thick part of the whitish Coloured whipworm and only a very small portion of the long, thin anterior part since the anterior part penetrates into the large intestinal wall for feeding purpose.




Fig 2: Magnified view showing clearly both the short, posterior thick part and also the anterior, thin part of the whipworm due to the higher magnification.

iv) Mild whipworm infection

Whipworm infection is clinically silent in the vast majority of cases, since the worm load tends to be low [13] and these patients require only anti worm treatment. Our study and also some other studies [3, 5, 13] have shown only a single whipworm in the colon while doing colonoscopy which represents the least load of whipworms and very mild whipworm infection in these fortunate patients.

v) Mild whipworm infection in our patient

Our patient was also found to have only mild whipworm infection with only a single whipworm in the colon while doing colonoscopy without anaemia (haemoglobin 14.4 g%) and without serious complications like Trichuris dysentery syndrome causing anaemia requiring intensive medical treatment and without colonic obstruction and perforation requiring surgical resection of the colon. The patient was treated with a single dose of 400mg of albendazole and were symptom-free after treatment with albendazole.

Conclusion

1. Whipworms are the most common nematodes or roundworms found in the large intestine of human beings while doing colonoscopy.
2. Whipworms do not suck blood and feed only on the tissue secretions of the large intestinal wall. Hence anaemia is not common in whipworm infection. Our patient also did not have any anaemia. In another study also, all the three patients with whipworm or trichuris trichiura found in colonoscopy did not have anaemia

3. Our study and also some other studies have shown only a single whipworm in the colon while doing colonoscopy which represents the least load of whipworms and mild whipworm infection in these fortunate patients.
4. Hence our patient was found to have only mild whipworm infection with only a single whipworm in the colon while doing colonoscopy without anaemia and without serious complications like Trichuris dysentery syndrome or colonic obstruction and perforation.

Acknowledgement

The author sincerely thanks the staff nurse Nithya who was assisting while doing colonoscopy and the staff nurses A.K. Selvi and Nithya for their immense help rendered to the author while conducting this work. The author acknowledges the immense help received from the scholars whose articles are cited and included in references of this manuscript. The author is also grateful to authors / editors / publishers of all those articles, journals and books from where the literature for this article has been reviewed and discussed.

References

1. Joo JH, Ryu KH, Lee YH, Park CW, Cho JY, Kim YS *et al.* Colonoscopic diagnosis of whipworm infection *Hepatogastroenterology.* 1998; 45(24):2105-9.
2. Do KR1, Cho YS, Kim HK, Hwang BH, Shin EJ, Jeong HB *et al.* Intestinal helminthic infections diagnosed by colonoscopy in a regional hospital during 2001-2008. *Korean J Parasitol.* 2010; 48(1):75-8.
3. Yoshida M, Kutsumi H, Ogawa M, Soga T, Nishimura

- K, Tomita S *et al.* A case of *Trichuris trichiura* infection diagnosed by colonoscopy. *Am J Gastroenterol.* 1996; 91(1):161-2.
4. Khuroo MS, Khuroo MS, Khuroo Trichuris NS. dysentery syndrome: a common cause of chronic iron deficiency anemia in adults in an endemic area (with videos). *Gastrointest Endosc.* 2010; 71(1):200-4.
 5. Ok KS1, Kim YS, Song JH, Lee JH, Ryu SH, Lee JH *et al.* *Trichuris trichiura* infection diagnosed by colonoscopy: case reports and review of literature. *Korean J Parasitol.* 2009; 47(3):275-80.
 6. Wang DD, Wang XL, Wang XL, Wang S, An Trichuriasis CL. diagnosed by colonoscopy: case report and review of the literature spanning 22 years in mainland China. *Int J Infect Dis.* 2013; 17(11):e1073-5.
 7. Tuan Sharif SE, Ewe Seng C, Mustaffa N, Mohd Shah NA, Mohamed Z. Chronic *Trichuris trichiura* Infection Presenting as Ileocecal Valve Swelling Mimicking Malignancy. *ISRN Gastroenterol.* 2011; 2011:105-178. doi: 10.5402/2011/105178. Epub 2010 Oct 31.
 8. Chang CW, Chang WH, Shih SC, Wang TE, Lin SC, Bair MJ. Accidental diagnosis of *Trichuris trichiura* by colonoscopy. *Gastrointest Endosc.* 2008; 68(1):154.
 9. Diniz-Santos DR, Jambeyro J, Mascarenhas RR, Silva LR. Massive *Trichuris trichiura* infection as a cause of chronic bloody diarrhea in a child. *J Trop Pediatr.* 2006; 52(1):66-8.
 10. Lorenzetti R1, Campo SM, Stella F, Hassan C, Zullo A, Morini S. An unusual endoscopic finding: *Trichuris trichiura*. Case report and review of the literature. *Dig Liver Dis.* 2003; 35(11):811-3.
 11. Tokmak N, Koc Z, Ulasan S, Koltas IS, Bal N. Computed tomographic findings of trichuriasis *World Journal of Gastroenterology*, 2006; 12(26):4270.
 12. Lee SH, Kwon JE, Cheong YS. Two cases of *Trichuris trichiura* infection diagnosed by colonoscopy. *Korean Journal of Family Medicine*, 2010; 31(8):622-629.
 13. Herman MA, Ukawa K, Sugawa C. CASE REPORT: Diagnosis and Removal of Cecal Whipworm Infection. *Digestive diseases and sciences*, 2000; 45(8):1639-1643.
 14. Azira MS, Zeehaida M. Severe chronic iron deficiency anaemia secondary to *Trichuris dysentery syndrome*-a case report. *Trop Biomed*, 2012; 29(4):626-631.